




CONTACT INFORMATION

Mining Records Curator
Arizona Geological Survey
416 W. Congress St., Suite 100
Tucson, Arizona 85701
602-771-1601
<http://www.azgs.az.gov>
inquiries@azgs.az.gov

The following file is part of the Doug K. Martin Mining Collection

ACCESS STATEMENT

These digitized collections are accessible for purposes of education and research. We have indicated what we know about copyright and rights of privacy, publicity, or trademark. Due to the nature of archival collections, we are not always able to identify this information. We are eager to hear from any rights owners, so that we may obtain accurate information. Upon request, we will remove material from public view while we address a rights issue.

CONSTRAINTS STATEMENT

The Arizona Geological Survey does not claim to control all rights for all materials in its collection. These rights include, but are not limited to: copyright, privacy rights, and cultural protection rights. The User hereby assumes all responsibility for obtaining any rights to use the material in excess of "fair use."

The Survey makes no intellectual property claims to the products created by individual authors in the manuscript collections, except when the author deeded those rights to the Survey or when those authors were employed by the State of Arizona and created intellectual products as a function of their official duties. The Survey does maintain property rights to the physical and digital representations of the works.

QUALITY STATEMENT

The Arizona Geological Survey is not responsible for the accuracy of the records, information, or opinions that may be contained in the files. The Survey collects, catalogs, and archives data on mineral properties regardless of its views of the veracity or accuracy of those data.

OTHER MINES AND PROSPECTS ON THE TENNESSEE VEIN

The Elkhart mine, at the extreme north end of the Tennessee vein, is an old mine that has been idle for many years. The total production from this mine from 1901 through 1948, as shown in table 2, has been small. The mine workings, now inaccessible, are reported to consist of three shafts, six levels (the lowest 500 feet deep), drifts totaling about 2,600 feet, numerous stopes, and several crosscuts.

The Silver Age mine, near the extreme south end of the vein, was primarily a silver mine (table 2). The silver was probably derived in large part from silver chloride (cerargyrite) found in the oxidized zone. Accurate data concerning the inaccessible mine workings could not be obtained. It is reported that the shaft is about 150 to 200 feet deep and that drifts and stopes extend northward from the shaft for some 200 or 300 feet. The vein material on the mine dump is partly oxidized, chiefly to iron hydroxides. Pyrite is the most abundant primary sulfide. Minor amounts of galena and sphalerite, together with sparse chalcopyrite, are associated with the pyrite in quartz gangue.

Several shafts have been driven and numerous pits and trenches have been dug along the Tennessee vein from the Silver Age shaft to the Tennessee shaft. The deepest of these is the Johnny Bull shaft (pl. 18), which is reported to be 88 feet deep. No drifting or stoping from this shaft is known.

Diamond drilling on the southern part of the Tennessee vein was carried out by the United States Bureau of Mines (Tainter, 1947) during the period from September 16 to December 8, 1943. The exploratory work consisted of eight drill holes on the Johnny Bull and Silver Knight claims, between 750 and 2,450 feet south of the Tennessee shaft. The holes were distributed along the vein at intervals ranging from 200 to about 375 feet. All holes were drilled from the surface and inclined toward the vein. Four were drilled from the west side of the vein outcroppings and the other four from the east side. Depths below the surface at which the vein was intersected ranged from about 100 to 350 feet, the deepest corresponding approximately in altitude to the 400-foot level in the Tennessee mine.

All holes intersected the vein, but the vein filling in seven of the eight cores was barren of ore minerals or was so low in grade as to be of little or no economic interest. The only hole that showed a substantial amount of the ore minerals was hole 8, located about 1,900 feet south of the Tennessee shaft. This hole intersected the vein about 100 feet below the surface, at an approximate altitude of 4,100 feet. A 3.5-foot interval of sphalerite, galena, and pyrite in quartz gangue assayed 7.6 percent zinc, 0.1 percent lead, and 0.03 percent copper.

This intersection might suggest that the top of an ore body was penetrated, but the Bureau of Mines engineers believed that the extensive drilling necessary to determine the existence of an ore shoot in the vicinity of hole 8 was not warranted.

TURQUOISE MINES

Deposits of turquoise are restricted to the Ithaca Peak granite and occur most abundantly in the southern half of the main intrusive body south of Mineral Park, particularly on Ithaca and Turquoise Peaks. Many small and shallow workings have explored these deposits, and only the larger ones are shown on plate 18. Some of the diggings are very old, having been started by the Aztec Indians. Very little work has been done on the deposits for many years.


Turquoise occurs typically in veinlets and small lenses in silicified, sericitized, and kaolinized porphyritic granite. Turquoise most commonly fills cavities in quartz veinlets, although some is in altered granite. Other minerals sparsely associated with turquoise in a few places are malachite, chrysocolla, and hydrous iron oxides. Sterrett (1908, pp. 847-852) describes some of the individual deposits in this area.

The features of the deposits suggest a secondary origin by supergene processes similar to those given by Paige (1912) for the origin of turquoise in the Burro Mountains of New Mexico.

LIST OF REFERENCES

The literature pertaining to the district is not extensive. The list given below includes the chief publications. Of these, Schrader's report on districts in Mohave County furnishes the most extensive description of the Wallapai district, and it is of particular value in furnishing descriptions of many of the mines. Thomas' manuscript contributes much information, particularly his detailed descriptions of the minerals and their paragenetic relationships. He includes a small-scale geologic map that covers an area extending from Mineral Park northwestward for several miles beyond Chloride. Most of the references are brief summaries of the geology and ore deposits, probably taken in part from Schrader's previous work.

- BASTIN, E. S., 1924, Origin of certain rich silver ores near Chloride and Kingman, Ariz.: U. S. Geol. Surv. Bull. 750, pp. 17-39.
 DARTON, N. H., 1925, A résumé of Arizona geology: Ariz. Bureau Mines Bull. 119, p. 180.
 DINGS, M. G., 1950, Wallapai mining district, Mohave County, Ariz.: Arizona Bur. Mines Bull. 156, pt. 1, pp. 138-142.
 ELSING, M. J., and HEINEMAN, E. S., 1936, Arizona metal production: Ariz. Bur. Mines Bull. 140, pp. 73-95.
 GARRETT, S. K., 1938, Tennessee-Schuykill mine: Ariz. Bur. Mines Bull. 145, pp. 117-119.



SCALE 1"